# **EI PASO HEALTH**

### **ROUTINE PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (12 YEARS- 20 YEARS)**

The following guideline provides recommendations for routine preventive services for children 12 years to 20 years. Children at increased risk may warrant additional services.

The following guideline provides recommendations for routine prevent									20
Recommendation	12 years	13 years	14 years	15 years	16 years	1/years	18 years	19 years	20 years
<b>History</b> : Documentation must contain an initial health history and each									
subsequent checkup must contain information on an interim history.	Х	Х	Х	Х	Х	Х	Х	Х	Х
Mental Health Screening: Mental health screening is required at each									
checkup and includes behavioral, social, and emotional development. One									
of the following validated, standardized mental health screening tools is	х	x	x	x	x	x	x	x	x
required when screening annually 12 through 18 years of age:		<u>I</u>		<u> </u>					
•Patient Health Questionnaire (PHQ-9, PHQ-A [anxiety, eating problems,									
mood problems and substance abuse])									
Patient Health Questionnaire Modified for Adolescents(PHQ-A									
[depression screen])									
Pediatric Symptom Checklist (PSC-17 and PSC-35)	Re	commende	d screening	g annually 1	L2 through	18 years of	age		
Pediatric Symptom Checklist Youth Report (Y-PSC)									
•Car, Relax, Alone, Forget, Family, and Trouble Questionnaire (CRAFFT)									
•Rapid Assessment for Adolesent Preventive Services (RAAPS)									
https://toolkits.solutions.aap.org/DocumentLibrary/BFTK2e_Links_Screeni									
ng_Tools.pdf			1	1	1		1		
Tuberculosis Screening: TB Questionnaire must be administered annually									
beginning at 12 months of age. A Tuberculin Skin Test is to be									
administered when the screening tool indicates a risk for possible									
exposure	х	х	х	х	х	х	х	х	х
Nutrition Screening: Dietary practices should be assessed to identify									
unusual eating habits such as pica, extended use of baby bottle feedings,									
or eating disorders in older children and adolescents. For nutritional									
problems, further assessment is indicated.	х	х	х	х	х	х	х	х	х
Age Appropriate Screening & Administration of Immunizations: Providers									
must assess the immunization status of clients at every medical checkup									
and vaccines must be administered according to the current Advisory									
Committee on Immunization Practices (ACIP). "Recommended Childhood									
and Adolescent Immunization Schedule.									
https://www.cdc.gov/vaccines/schedules/	х	х	х	х	х	х	х	х	х

EPHP4012109 QIC Approved 07/27/2021

# **EI PASO HEALTH**

### **ROUTINE PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (12 YEARS- 20 YEARS)**

The following guideline provides recommendations for routine preventive services for children 12 years to 20 years. Children at increased risk may warrant additional services.

12 years	13 years	14 years	15 years	16 years	17years	18 years	19 years	20 years
dia, dys,	dia, dys,	dia, dys,	dia, dys,	dia, dys,	dia, dys,	dia, dys,	dia, dys,	dia, dys,
std, hiv	std, hiv	std, hiv	std, hiv	std, hiv	std, hiv	std, hiv	std, hiv	std, hiv
				hiv (Once at	16 - 18 years)	dys (Once at 18 - 20 years)		
х	х	х	х	х	х	х	х	х
5								
х	х	х	х	х	х	х	х	х
х	х	х	х	х	х	х	х	х
х	х	х	х	Х	х	х	Х	х
х	х	х	х	х	х	х	х	х
	dia, dys, std, hiv  X S X X	dia, dys, std, hiv std, hiv  X X X X X X X	dia, dys, std, hiv std, hiv std, hiv  X X X X  S X X X X  X X X  X X X	dia, dys, std, hiv st	dia, dys, std, hiv std, hiv std, hiv hiv (Once at X X X X X X X X X X X X X X X X X X	dia, dys, std, hiv std, hiv std, hiv std, hiv hiv (Once at 16 - 18 years)  X X X X X X X X X X X X X X X X X X X	dia, dys, std, hiv st	dia, dys, std, hiv st

<sup>•</sup>Use the World Health Organization (WHO) growth charts http://www.cdc.gov/growthcharts/who\_charts.htm for infants and children birth to 2 years of age.

<sup>•</sup>Use the Centers for Disease Control and Prevention (CDC) growth charts http://www.cdc.gov/growthcharts/cdc\_charts.htm for children who are 2 years of age or older.

Vision: Visual Acuity results	х			х			х		
Subjective vision		х	х		х	х		x	х
Hearing Audiometric Screening results				х					
Subjective hearing	Х	Х	Х		Х	Х	Х	Х	Х

EPHP4012109 QIC Approved 07/27/2021

# **EI PASO HEALTH**

#### ROUTINE PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (12 YEARS- 20 YEARS)

The following guideline provides recommendations for routine preventive services for children 12 years to 20 years. Children at increased risk may warrant additional services.

Recommendation	12 years	13 years	14 years	15 years	16 years	17years	18 years	19 years	20 years
Age Appropriate Anticipatory Guidance & Health Education: Health									
education is designed to help parents and caregivers understand what to									
expect in terms of the child's development and to provide information for									
all ages about the benefits of healthy lifestyles and practices, as well as									
accident and disease prevention. Diet and Exercise are topics that are									
strongly encouraged.									
https://hhs.texas.gov/doing-business-hhs/provider-portals/health-services-	-								
providers/texas-health-steps/medical-providers	Х	Х	Х	Х	Х	Х	Х	Х	Х
<b>Dental:</b> Limited oral screening for caries and general health of the teeth									
and oral mucosa is part of the physical examination. In addition to the									
federal requirements, Texas Health Steps policy requires referral to a									
dentist at six months of age and every six months thereafter until the									
dental home has been established.	х	х	х	Х	х	Х	Х	Х	Х
Follow up Instructions to Return for Next Preventive Visit:	х	х	х	х	х	х	х	х	х

This guideline is based on the Texas Health StepsPeriodicity Schedule and the Texas Health Steps Checkup Components. https://hhs.texas.gov/doing-business-hhs/provider-portals/health-services-providers/texas-health-steps/medical-providers. Individual patient considerations and advances in medical science may supercede or modify these recommendations.

Revised June 1, 2021

EPHP4012109 QIC Approved 07/27/2021